

# **Patterns of Utilization of Maternal Healthcare Services in Haryana**

**Sonal Katyal**

**Assistant Professor (Economics)**

**GD Goenka World Institute | Lancaster University**

**Gurgaon, Haryana**

## **Introduction**

One of the eight Millennium Development Goals (MDGs) set by United Nations in 2000 is to improve the maternal health targeting at reducing the maternal mortality ratio by three quarters between 1990 and 2015 and to achieve universal access to reproductive health by 2015 . Maternal mortality is unacceptably high about 800 women die per lakh live births from pregnancy or child birth related complications around the world every day. In 2013, 2, 89,000 women died during and after pregnancy and childbirth. Ninety nine percent of these deaths occur in developing countries (WHO, 2014)

India has taken several steps in this regard since independence in the form of five year plans to promote the goal of universal health. But finally in 1983 it came up with a National Health Policy which aimed at Health for all by 2000 through universal comprehensive primary health services. But it failed to achieve it because of lack of financial resources and public health administrative capacity. In 2002, it attempted a new policy framework as the National Health Policy 2002 which primarily aimed at achievement of health goals keeping in mind the prevailing socio-economic conditions (India Health Report , 2003) .

Though National Health Policy has set objectives keeping in mind everyone's health needs but maternal health care needs a special attention as women are considered as the most vulnerable group in society.

Health and socio-economic development of any country are interdependent. Levels of maternal mortality tell us about the risk attributable to pregnancy and child birth as well as the performance of health systems in terms of access to health care and quality

of care provided. Yet the economic development in India has gained momentum over the last decade, but the public health system is lagging behind. In Haryana, rural population accounts for 71 percent of the state's population. Haryana's per capita income is amongst the highest and fastest growing in the country (Haryana.gov.in). But the Anthropological Survey of India reports that the position of women in Haryana continues to be bad. Women have lagged behind in all the sectors. Haryana is the second richest state in India in per capita income but sex ratio in Haryana continues to be highly unfavorable to females.

### **Significance of Study**

The study of maternal health care use in Haryana is timely because maternal mortality and morbidity still remain high in this state in spite of all the efforts made to address the healthcare needs of women of reproductive age. The use of National Family Health Survey-III data allows for a comprehensive analysis of maternal healthcare usage. This study is critical in enhancing the quality and effectiveness of national programs for maternal and child health. The main motivation behind this study is to explore the health system in terms of access. Secondly, not many studies have focused on Haryana State per se and not much literature is available despite unfavorable sex ratio. Hence, this topic is worth to be taken in consideration. This study provides an analysis for antenatal care utilization in Haryana and fills a critical gap in understanding this public health issue in this state and will help in future planning and implementation of such maternal health programmes which will aim at improving the health system.

### **Objective of the study**

The main objective of this study is to examine the differentials in utilization of maternal healthcare service i.e. antenatal services in Haryana on the basis of various socioeconomic and Sociodemographic factors.

Anderson's Behavioral Theory (Anderson, 1968, Anderson and Newman, 1973) is used as a conceptual framework for the analysis of maternal health care use based on

the socio-economic characteristics as potential correlates. The Anderson behavioral model postulates that the use of maternal healthcare services is influenced by three sets of individual characteristics (a) Predisposing characteristics (i.e. age, parity, woman's education, her husband's education, woman's autonomy); (b) Enabling characteristics (i.e. household wealth , place of residence , woman's employment) ; (c) Need characteristics (i.e. antenatal care, delivery care , postnatal care).

Considering the Anderson's Behavioral theory model predisposing factors are socio demographic that would predispose women to use maternal healthcare services, while the enabling factors are the socioeconomic factors that may make women prove to use healthcare services

In this study the Aday and Anderson's model was not tested rather the association between explanatory variables (demographic and social) and utilization of maternal healthcare indicators were explored. Factors affecting maternal healthcare utilization are based on this framework but the variables used in this study are identified after reviewing literature and keeping in mind the background characteristics of population.

## **Review of Literature**

Determinants or correlates of utilization can emanate from demand or supply side.

### **Demand side Determinants**

Demand side determinants are individual and household level factors that affect the utilization of services.

The household socio-economic status and mother's education were the most important factors associated with the use of Antenatal Care and skilled attendance at delivery in Madhya Pradesh. (Jat, Ng, & Sebastian, 2011)

Women's age at child birth appears to be a related factor to the use of maternal healthcare services. In a study based on India (Bhatia and Cleland, 1995) found that mothers younger than 18 were less likely to use antenatal services compared to women above 18 years of age. While some studies such as (Chakraborty et al., 2002)

suggests that women's age can also indicate women's knowledge of healthcare services and can have a positive effect on utilization of maternal healthcare services

A study based on north-eastern states of India suggests that for better maternal health care utilization, policies should be directed towards the vulnerable group i.e. women representing the SC/ST background because high percentage of population belongs to scheduled tribes in the north-eastern states. The result showed that education of both the woman and her husband plays a vital role in utilization of maternal health care services. Other variables that can have significant impact on availing of the maternal health care services are media, autonomy enjoyed by women and their occupational status (Chakrabarti and Chaudhuri, 2007).

A study based on Uttar Pradesh suggests that women's autonomy plays an equally important role in maternal health care utilization as women's educational status. The analyses show that women with greater freedom of movement are more likely to use safe delivery care and obtain higher levels of antenatal care. (Bloom, Wypij, & Dasgupta, 2001).

The utilization of maternal healthcare services also depends on geographical and topographical characteristics of a state. The number of women utilizing antenatal services is less in Uttaranchal which is a hilly area in comparison to Jharkhand and Chhattisgarh (Pandey, Roy, Sahu and Acharya, 2004). According to the study (Bhattacharya and Tandon, 1991), mother's educational level has a positive impact on utilization of maternal healthcare services. Visit of a health worker has a significant positive impact on the utilization of full antenatal and post natal care services among women (Sunil et al., 2006)

Bhatia and Cleland (1995) suggested that for improving maternal health services women should have adequate knowledge of health-care seeking behavior during the reproductive process and its determinants.

Not only women's education, a study shows that husband's education has a significant influence on the adequacy of maternal healthcare utilization in India. Maternal care usage was measured by a composite index based on responses of both

antenatal and delivery care and categorized from poor to excellent and husband's education was found to be linked with excellent maternal healthcare use (Sunil et al, 2006).

A study conducted in rural North India suggests that there is a caste divide as majority of traditional birth attendants called Dias belong to the lower caste and trained birth attendants such as nurses or doctors belong to upper caste. For providing maternal healthcare services both the caste groups have to be in contact with each other, upper caste health providers prefer to cater to upper caste women and lower caste women choose to utilize services offered by traditional birth attendants for the sake of avoiding embarrassment of caste discrimination (Saroja, Altarac & Sibley, 2008).

Pallikadavath et al. (2004) found that Muslim women utilized these services more than others whereas, there was low utilization of services among scheduled castes and scheduled tribes. A study observed that Muslim women give birth outside medical facilities more often than Hindu women. But nothing has clearly emerged in India on utilization of healthcare services based on religion.

Another study found that women who have a job that paid them in cash were more likely to utilize healthcare services during their pregnancy (Chakraborty et al., 2002).

Other factors such as economic status of women of the household also affect the utilization of antenatal care services and delivery care (Pandey et al., 2002). A study was conducted on Bihar, Madhya Pradesh, Rajasthan, and Uttar Pradesh using National Family Health Survey (1998-99) found that higher social and economic status is associated with increased utilization of antenatal care including blood pressure measurement, blood test and urine test, but the obstetric physical examination depends on education of women and their husbands. Women from poor and uneducated backgrounds with at least one child are less likely to receive antenatal care services (Pallikadavath et al., 2004)

### **Supply Side Determinants**

Utilization of healthcare services is possible only if healthcare services are available for access. Healthcare facilities in terms of proximity and providers of healthcare play an important role in utilization of services.

A study based on Andhra Pradesh, Karnataka, Kerala and Tamil Nadu found that variations in use of services were primarily related to availability and access (Navaneetham & Dharmalingam, 2002).

Utilization of healthcare services also depends on access to health facility. However, (Bhattacharya and Tandon, 1991) found out that even if rural women are staying near health centre still they do not utilize healthcare services.

Women who lived near a village health worker/nurse were more likely to receive adequate and early antenatal care visits than women without a village health worker (Nielsen et al., 2001).

Another study based on Nepal has shown that services are utilized twice as high if health facility is in the community and also if there are regular visits by health workers (Acharya and Cleland, 2000).

This literature review has highlighted the importance of individual, household and community level factors that affect the use of antenatal services.

### **Hypothesis**

Based on literature review and previous empirical research, the following associations are expected. Younger mothers (less than 18 years of age) have lower odds of using antenatal care than mothers above 18 years of age. Higher level of education for both the woman and her husband will be positively associated with antenatal care usage. Women who have high autonomy (decision making power and mobility) are more likely to utilize these maternal healthcare services compared to women with low autonomy. Women with any mass media exposure (read magazine, watch Television and listen to radio) will utilize these maternal healthcare services more than women with no mass media exposure. Women with lower levels of household wealth will be less likely to use antenatal care services. Rural residence will be negatively associated

with antenatal care usage. Women who have a job are more likely to have adequate antenatal care. Women who have met with an auxiliary nurse midwife (ANM) or community health worker are more likely to utilize maternal healthcare services. Women belonging to other castes will utilize these services more. Hindu women are more likely to use these services.

## **Methods**

### **Data Source**

The present study uses the third round of the National Family Health Survey (NFHS) data which is similar to the Demographic and Health Surveys (DHS). DHS collects, disseminates national data on health and population in developing countries.

The NFHS is a large scale, multi round survey conducted in a representative sample of households throughout India. The NFHS-III was conducted in 2005-06 and is the third in the series of surveys. All three NFHS surveys were conducted by the Ministry of Health and Family Welfare (MOHFW), Government of India. Indian Institute of Population Sciences was selected as the nodal agency for carrying out the survey. NFHS-III covers information on nationally representative sample of 1, 09,041 Households, 1, 24,385 (both married and unmarried) Women age 15-49 years and 74,369 Men (both married and unmarried) age 15-54 years from all 29 states.

### **Sampling Plan**

The sampling plan and design explained in this study is done by International Institute of Population Sciences (IIPS), Mumbai who conducted NFHS –III survey.

NFHS-III adopted a two stage sample design in most rural areas and three stage sample design in most urban areas in all the 29 states. In each state, the rural sample was typically selected in two stages: the first stage involved selected of primary sampling units (PSU's), which are villages with probability proportional to population

size (PPS), the second stage involved the systematic selection of households within each PSU.

Stratified sampling was adopted. In rural areas, list of villages was taken from 2001 census and it served as the sampling frame. Stratification was done to ensure the inclusion of villages with different socio economic characteristics in the sample. The first level of stratification was geographic, with districts being subdivided into contiguous regions. The selection of required number of households from the list of households was done using systematic sample. The overall response rate for women was 92.4 percent whereas for men it was 84.9 percent.

### **Study Population and Sample Size**

The present study examines the pattern of utilization of maternal health services among currently married women age (15-49) years in Haryana.

In Haryana, NFHS-III is based on a sample of 2,302 households that is representative at the state level and within state at the urban and rural levels. The study interviewed 2,790 women age 15-49 years from all sample households and 1,083 men age 15-54 from a sub sample of households for obtaining information related to population, health and nutrition in the state. The household response rate in the state as a whole was 99 percent and the individual response rates were 96 percent for eligible women and 85 percent for eligible men.

In this study data related to currently married women (15-49) years of age is examined and a total of 2134 currently married women from both rural and urban areas is included.

### **Measurement**

#### **Dependent Variables**

The study measures the outcome variables namely Antenatal care to explore inequities in utilization pattern of maternal healthcare services in Haryana. The selected indicator of maternal healthcare utilization and its components are examined



on the basis of guidelines laid by the Ministry of Health and Family Welfare (MOHFW), Government of India and World Health Organization (WHO).

For this study, Antenatal care includes those women who had a minimum of three antenatal care visits. The MOHFW recommends that a woman should have at least three antenatal care visits during pregnancy and this standard was used to construct this dependent variable. In this study, **Antenatal care** was coded as a binary dependent variable, with a value of 1 if the woman had at least three antenatal visits and a value 0 if the woman had less than three antenatal care visits.

### **Independent Variables**

The independent variables used in this study are socio-economic and socio-demographic.

### **Predisposing Factors**

**Women's age at child birth** is defined as women's age in years. A variable was created (womanaage). Womanaage was coded as 0 if **women's age** was less than 18 years and was coded as 1 if women's age is greater than 18 years.

**Women's and their Husband's education level** are defined using highest education level. Variables Womeneduc and Husbandeduc were created and were categorized into 0,1,2,3. It was coded as 0 if they had no education, coded as 1 if the woman had studied till primary, coded as 2 if till secondary and coded as 3 if till higher level of education.

**Religion** of the woman was categorized as Hindu, Muslim, and Sikh. Religion was coded as 0 if Hindu, coded as 1 if Muslim and coded as 2 if Sikh.

**Caste** is defined according to the social group that woman belongs to as per the categorization and identification done in NFHS-3 data. Variable caste is coded as 0 if women belong to none of them, coded as 1 if women belong to scheduled castes (SCs), coded as 2 if women belong to scheduled tribes (STs) and coded as 3 if women belong to other backward classes (OBCs).

**Women's Autonomy** was computed by taking into account decision making and women's mobility. Decision making was computed on the basis of decision about healthcare for yourself, decision about making major household purchases, decision about making purchases for daily household and decision about visits to your family or relatives. These four factors were combined and were coded as 0 if respondent and respondent plus husband take decisions and were coded 1 if someone else is taking decision on respondent's behalf. Women's mobility was computed by taking into account following factors whether women is allowed to go to market, allowed to go to health facility, allowed to visit places outside the village/community. These three factors were combined and were coded as 0 if women were allowed to go alone or with someone else and were coded as 1 if they were allowed to go anywhere. Both decision making and women mobility were combined to construct a variable wautonomy using information from NFHS-3 data. Wautonomy was coded as 0 if women were allowed to go out and were involved in decision making and was coded as 1 otherwise.

**Mass media exposure** was computed by taking into account how often respondents read magazine, listen to radio and watch television. They were coded as 0 if they don't read magazine, listen to radio and watch television and coded as 1 otherwise. All three factors were combined to create variable media and then from media we created media exposure and coded it as 0 if they had no exposure and coded as 1 if they had some exposure.

### **Enabling Factors**

**Place of residence** is defined as the place of stay. It is coded as zero if the woman stays in urban area and is coded as 1 if woman stays in rural area.

**Wealth index** is defined as woman falls into which category (poorest, poorer, middle, richer and richest) and this categorization is taken from NFHS-3 data. Poorest is coded as 0, poorer as 1, middle as 2, richer as 3 and richest as 4.

**Met with ANM/Community health worker** is defined as whether in last 3 months woman has met any ANM/Community health worker. It is coded as 0 if not met and coded as 1 if otherwise.

**Work status** is defined whether the woman is working or not. If the woman is not working it is coded as 0 and 1 otherwise.

### **Statistical Methods**

Univariate analytic techniques were used to provide a description of all variables used in the analysis. The NFHS -3 dataset was used to extract information and STATA 10 was used to correctly estimate population means, proportions and standard deviations.

To identify factors associated with maternal healthcare utilization, bivariate and multivariate analyses were performed. Bi-variate analyses were performed to examine the nature of association between utilization of maternal healthcare services by selected socioeconomic and demographic background characteristics.

Multivariate analyses were performed to examine the nature of predictor variables and their relative contribution in explaining the dependent variables. Since both outcomes in this analysis are dichotomous in nature, logistic regression was used to predict the association between the dependent variables and the independent variables. Ordinary least square regression would be inappropriate as it assumes error variances (residuals) to be normally distributed. The binary response(y) for each individual was related to a set of categorical predictors (X), logit of the probability of each outcome can be modeled as:

$$\text{Logit}(\pi_i) = \log(\pi_i / 1 - \pi_i) = \beta_0 + \beta(X) + \varepsilon$$

The probability of an individual who had received antenatal care or had safe delivery or received assistance in postnatal care is  $\pi_i$ .  $\beta_0$  estimates the log odds of antenatal care or had safe delivery or received assistance in postnatal care for the reference group, and the parameter  $\beta$  estimates with maximum likelihood, the differential log odds of full antenatal care or had safe delivery or received assistance in postnatal care are associated with the predictor X, in comparison to reference group. E represents the

error term. Odds ratio and confidence intervals for each independent variable were presented for the interpretation of the coefficients. This analysis was performed using STATA10.

Before running the analysis, the models were checked for multicollinearity using the variance inflation factor (VIF). The VIF measures how much of the variance of the coefficient estimate is being by multicollinearity. VIF values greater than 10 indicate multicollinearity. The models used in this study satisfy this condition and this indicates that covariates are not collinear. Therefore, no explanatory variable was dropped from the analysis.

## Results

### Profile of the Respondents

Table 1 represents the percentage of women (15-49) years of age who had at least one live birth during the last five years preceding the survey by selected background characteristics. Majority of women were above 18 years of years. With regard to education, 46% of women had no education but 57% of Husband's were educated till secondary level. Most of the women 88% were Hindu. As per the social group most women were from other backward classes (OBCs).around 63% women enjoy high autonomy and 74% have exposure to any of the mass media. Most of the women are not working and residing in rural areas. 24% and 29% of women fall in middle and richer wealth index respectively. 92% of women had not met any health worker in the past three months.

**Table 1 Percentage of women (15-49) years who had at least one live birth during the last five years preceding the survey by background characteristics, NFHS-3(2005-06)-Haryana**

<b>Background characteristics</b>	<b>N (Number)</b>	<b>%(Percentage)</b>
<b>Woman's age</b>		
Less than 18 years	31	1.45
Greater than 18 years	2,103	98.55
<b>Woman's education</b>		

No Education	966	45.27
Primary	278	13.03
Secondary	758	35.52
Higher	132	6.19
<b>Husband's education</b>		
No Education	442	20.74
Primary	223	10.46
Secondary	1,225	57.48
Higher	241	11.31
<b>Religion</b>		
Hindu	1891	88.78
Muslim	125	5.87
Sikh	114	5.35
<b>Caste</b>		
Others	1,181	55.39
Scheduled Castes(SC's)	488	22.89
Scheduled Tribes( ST's)	14	0.66
Other Backward Classes (OBCs)	449	21.06
<b>Background characteristics</b>	<b>N (Number)</b>	<b>%(Percentage)</b>
<b>Work Status</b>		
Non- Working	1646	77.13
Working	488	22.87
<b>Autonomy</b>		
Low	794	37.21
High	1,340	62.79
<b>Media Exposure</b>		
No exposure	557	26.10
Any exposure	1,577	73.90
<b>Wealth Index</b>		
Poorest	72	3.37

Poorer	243	11.39
Middle	522	24.46
Richer	611	28.63
Richest	686	32.15
<b>Place of residence</b>		
Urban	553	25.91
Rural	1,581	74.09
<b>Met with health worker in last 3 months</b>		
No	1,948	91.28
Yes	186	8.72

Note: all N are unweighted.

### **Differentials in Utilization Pattern of Antenatal care Services**

This section identifies the factors associated with the utilization of antenatal care, services; we examined the bi-variate differential of the selected socio economic and demographic characteristics. This was done by finding the association between predisposing and enabling factors and the use of antenatal services. Chi- square statistics are used for differences in distribution of predisposing and enabling factors among women (15-49) years of age who had at least one live birth in last five years utilizing antenatal care services

Table 2 shows the percentage of women who utilized antenatal care services by selected background characteristics. The pattern of utilization of antenatal care is high for women who are above 18 years of age. Women who are not working and enjoy more autonomy and have any mass media exposure utilize antenatal care services more. On the other hand, women with no education utilize these services more. Rural women who have not met any health worker in past 3 months take more antenatal

care. Women belonging to other social groups and richest wealth quintile utilize these services more than women belonging to other castes and lower wealth quintiles.

**Table 2 Percentage of women (15-49) years who had at least one live birth during the last five years preceding the survey by antenatal care utilization, NFHS-3(2005-06)-Haryana**

<b>Background characteristics</b>	<b>%(Percentage)</b>
<b>Woman's age</b>	(0.0005) <sup>ns</sup>
Less than 18 years	1.45
Greater than 18 years	98.55
<b>Woman's education</b>	(19.5009) <sup>***</sup>
No Education	43.45
Primary	13.33
Secondary	36.33
Higher	6.86
<b>Husband's education</b>	(41.9035) <sup>***</sup>
No Education	18.72
Primary	9.78
Secondary	59.27
Higher	12.23
<b>Religion</b>	(130.1645) <sup>***</sup>
Hindu	90.83
Muslim	3.35
Sikh	5.81
<b>Caste</b>	(5.4949) <sup>ns</sup>
Others	56.47
Scheduled Castes(SC's)	22.27
Scheduled Tribes( ST's)	0.61
Other Backward Classes (OBCs)	20.65
<b>Work Status</b>	(0.4906) <sup>ns</sup>
Non- Working	76.85
Working	23.15



<b>Background characteristics</b>	<b>%(Percentage)</b>
<b>Autonomy</b>	(16.1469)***
Low	39.04
High	60.96
<b>Media Exposure</b>	(48.9195)***
No exposure	23.20
Any exposure	76.80
<b>Wealth Index</b>	(113.5240)***
Poorest	2.40
Poorer	9.54
Middle	23.20
Richer	29.39
Richest	35.47
<b>Place of residence</b>	(22.7386)***
Urban	22.89
Rural	72.11
<b>Met with health worker in last 3 months</b>	(11.6243)***
No	92.19
Yes	7.81

Note: Figures in parentheses are chi-square statistics; this test was applied for each variable

Level of significance: \*p<.10; \*\*p<.05;\*\*\*p<.01; ns: not significant

### **Determinants of Antenatal care Utilization**

Table 3 shows the results of multivariate analysis of antenatal care among women (15-49) years of age who had at least one live birth during the past five years. The significant determinants for antenatal care used in the analysis are woman's age at child birth, woman's education level, her husband's education level, mass media exposure, autonomy, wealth, place of residence, meeting with a health worker, work status, religion and caste.

Women less than 18 years of age, women with no education, husband's with no education, women belonging to Hindu religion, women belonging to other social groups, women who are not working, women who have low autonomy, women who have no mass media exposure, women belonging to poorest wealth index, women residing in urban areas and women who have not met any health worker in past 3 months are taken as reference category.

Women who are not adolescents i.e. above 18 years of age tend to utilize antenatal care services two times (OR = 2.960652 ; CI=.8845427 – 9.909596) more than adolescent women i.e. below 18 years of age. Woman's education level primary, secondary or higher did not show any significant increase in utilization of antenatal services in comparison to woman with no education. The odds of receiving antenatal care were high for women whose husband's had secondary or higher level of education in comparison to women whose husband had no formal education. There was no significant difference in utilization of antenatal care services for Muslim women but odds for receiving services were higher for Sikh women (OR =1.491157; CI=.7480687 – 2.972388) . The odds of receiving antenatal care were highest for women belonging to other backward classes (OR =1.066138; CI=.655831 – 1.292942) in comparison to women belonging to scheduled castes (OR = 1.019197; CI= .7392245 – 1.405206) or scheduled tribes (OR = 0.9802248; CI=.2494888- 3.851238). Work status of women had positive effect on utilization of antenatal care services (OR = 1.334586; CI=.9841091 – 1.809882)

Woman's autonomy did not have any significant effect on utilization of antenatal care services.

The odds ratio indicates that women who had any exposure (OR = 1.194362; CI=.8680895 – 1.643277) utilize these services more than women with no exposure. Wealth index showed a notable effect on the utilization of antenatal care services. Women from richer and richest index are nearly two (OR = 2.417466; CI=1.246266 – 4.689323) and four (OR=4.950028CI=2.355501- 10.40237) times more likely to utilize these services respectively compared to women belonging to poorer (OR = 1.019907; CI=.5428392 – 1.916241) and middle (OR = 1.530569; CI=.8242293 –

2.842221) wealth quintile. Women residing in rural areas did not show any significant increase (OR = .8451361; CI=.5882021-1.214302) in utilization of services than women staying in urban areas. Women who have met with a health worker in last three months did not suggest any increase in utilization of antenatal care services in comparison to women who have met with a health worker in last three months.

The analysis suggests that significant factors affecting utilization of antenatal care services among women 15-49 years of age who had at least one live birth in last five years are woman's age, wealth index, husband's education, media exposure, woman's work status, caste and religion.

**Table 3 Logistic Regression showing odds ratio and 95 % confidence interval for receiving antenatal care among women (15-49) years who had at least one live birth during the last five years preceding the survey, NFHS-3(2005-06)-Haryana**

<b>Covariates</b>	<b>Odds ratio</b>	<b>95 % CI</b>
<b>Woman's age</b>		
Less than 18 years*	1.000	
Greater than 18 years	2.960652	.8845427 – 9.909596
<b>Woman's education</b>		
No Education*	1.000	
Primary	.7815156	.5120718-1.192736
Secondary	.515813	.3600605-.7389399
Higher	.5653594	.2354061-1.357787
<b>Husband's education</b>		
No Education*	1.000	
Primary	0.9005271	.5834772- 1.389855
Secondary	1.220119	.8529871 – 1.745267
Higher	1.199789	.6283673 – 2.290846

<b>Covariates</b>	<b>Odds ratio</b>	<b>95 % CI</b>
<b>Religion</b>		
Hindu*	1.000	
Muslim	.1879701	.1186357 - .2978257
Sikh	1.491157	.7480687 – 2.972388
<b>Caste</b>		
Others*	1.000	
Scheduled Castes(SC's)	1.019197	.7392245 – 1.405206
Scheduled Tribes( ST's)	0.9802248	.2494888- 3.851238
Other Backward Classes (OBCs)	1.066138	.7659536 – 1.483967
<b>Work Status</b>		
Non- Working*	1.000	
Working	1.334586	.9841091 – 1.809882
<b>Autonomy</b>		
Low*	1.000	
High	.5260842	.3962889 - .8983909
<b>Media Exposure</b>		
No exposure*	1.000	
Any exposure	1.194362	.8680895 – 1.643277
<b>Wealth Index</b>		
Poorest*	1.000	
Poorer	1.019907	.5428392 – 1.916241
Middle	1.530569	.8242293 – 2.842221
Richer	2.417466	1.246266 – 4.689323
Richest	4.950028	2.355501- 10.40237
<b>Place of residence</b>		
Urban*	1.000	
Rural	.8451361	.5882021-1.214302
<b>Met with health worker in last 3 months</b>		
No*	1.000	
Yes	.6712552	.4494421 – 1.00254

Note: \* = reference category

## **Discussion & Conclusion**

This study examined differentials in the use of maternal health care services namely antenatal care in Haryana on the basis of various socio – economic factors such as woman's age at child birth, woman's education level, her husband's education level, mass media exposure, autonomy, wealth, place of residence, meeting with a health worker, work status, religion and caste. The relationship between outcome variable and predisposing and enabling factors is based on Anderson's model. The data was analyzed to find the relationship between predisposing and enabling factors and the outcome.

Bivariate analysis of antenatal care utilization with predisposing and enabling factors show several patterns.

Women who are not adolescents i.e. above 18 years of age tend to utilize antenatal care services two times more than adolescent women i.e. below 18 years of age. A study based in rural north India (Pallikadavath et al., 2004) also explain that age at marriage was positively associated with access or attendance for ANC. Antenatal check-ups were more likely among women whose age at the time of marriage was 19 years or above as compared to those women who married at younger age.

The odds of receiving antenatal care were high for women whose husband's had secondary or higher level of education in comparison to women whose husband had no formal education. This also conforms to another study that shows Husband's education was a statistically significant predictor in Andhra Pradesh but not in Karnataka (Navaneetham and Dharmalingam, 2002). The odds of receiving antenatal care were highest for women belonging to other backward classes in comparison to women belonging to scheduled castes or scheduled tribes. The low utilization of maternal health care among certain groups shows the lack of access to health care services among socially backward communities. This was demonstrated by another study based on rural Hindu women in Maitha, Uttar Pradesh which suggests that majority of Dais, untrained TBA's belong to the lower caste and trained TBA's belong to upper caste. Because of class discrimination and to avoid physical contact at

the time of service with lower caste women Trained TBA's only cater to upper caste women (Saroaha, Altarac, Sibley, 2008)

Women from richer and richest index are nearly two and four times more likely to utilize these services respectively compared to women belonging to poorer and middle wealth quintile. Household economic status has a positive impact on use of ANC. This is proved by a study that women with high economic status were more likely to receive adequate and early ANC than those with low economic status (Magadi et al. 2000, Matsumura & Gubhaju 2001).

### **Limitations of the study**

The study had some limitations, which need to be considered while interpreting the results.

1. The study used the data from a cross sectional survey and the association between explanatory variables and the indicator of the use of maternal health services was examined, could not draw conclusions about causality.
2. Some correlates of maternal health care utilization are missing from our analysis such as distance of health facilities from the locality of residence, and this could have influenced the patterns of utilization of maternal health services. Since data on this variable was not available in the survey.

### **Recommendations**

Efforts need to be made at community as well as household level to improve maternal health care services.

At community level, there should be call for action to reduce financial barriers while increasing awareness about the maternal healthcare services, particularly among women belonging to low wealth quintile. Efforts should be put to educate the men about the importance of maternal health care services. There should be more schemes for scheduled castes, scheduled tribes and other backward classes. Not just the literacy, there should be overall development of women. They should be exposed to newspapers, radio, television etc. the most important indicator of maternal health care is antenatal care as delivery care and post natal care are linked to this indicator. So,

more emphasis should be laid on improvement of antenatal care services. At Household level, Husband and mother in law should be used as targets for messages. There should be pregnant women groups and some incentive based competition among mothers to be model mother who can set an example for other mothers.

Health is a social phenomenon whose determinants cannot be separated from other social and economic determinants. Haryana has progressed at some fronts but still large gaps persist between the need and provision of services. It will require monitoring and evaluation of ongoing programs and effective implementation.

Health services no matter how efficient cannot change the condition of the marginalized people unless they are helped to become self-reliant and the root problems are addressed. People who are poor and illiterate are like uncut gems hidden under the dirt and stone. Given the opportunity, they can reach their full potential and live as responsible, sensitive human beings, possessing self – reliance and the liberty to shed those old customs and traditions that impede health and development (Arole and Arole, 1994)

### **Future Scope of Study**

Further on, the other components of maternal healthcare services such as delivery care and postnatal care can be examined based on the various socioeconomic and Sociodemographic variables.

## **Bibliography**

- Acharya, L.B. & Cleland, J. (2000). Maternal and child health services in rural Nepal: does access or quality matter more? *Health Policy and Planning* 15(2), 223-229.
- Aday, Lu, Ann & Anderson, Ronald M. (1981). Equity of access to medical care: A conceptual and empirical overview: *Medical Care*, 19(12), Supplement.
- Anderson, Ronald; Aday, Lu Ann (1978). Access to medical care in the US: Realized National Health Policy (2002) [http://www.iapsmhc.org/userfiles/National\\_Health\\_policy\\_2002-2.pdf](http://www.iapsmhc.org/userfiles/National_Health_policy_2002-2.pdf) as accessed Feb 2014
- Arole and Arole (1994) *Jhamkhed: A Comprehensive Rural Project*, Macmillan, 1<sup>st</sup> Edition .
- Bhatia, Jagdish. & Cleland , John. (1995). Determinants of maternal care in regions of South India . *Health Transition Review* 5, 127-142
- Bhattacharya , R. & Tandon, J. (1991) . Managerial Gaps in the Delivery of Antenatal Care Services in a Rural Area of Varanasi. *Indian Journal of Public Health*, 35(4), 113-116
- Bloom, Shelah S.; Wypij, David and Gupta D Monica (2001). Dimensions of women's autonomy and the influence on maternal health care utilization in a north Indian city: *Demography*, 38(1), 67-78
- Chakrabarti, Anindita & Chaudhari, Kaushik (2007). Antenatal and maternal health care utilization: Evidence from north-eastern states of India: *Applied Economics*, 39, 683-695.
- Chakraborty, N., Islam, M.A., Chowdhury, R.I., & Bari, W. (2002). Utilisation of postnatal care in Bangladesh: evidence from a longitudinal study. *Health & Social Care in the Community* 10(6), 492-502.
- EconomicSurveyofHaryana<http://esaharyana.gov.in/Data/Economic%20Survey%20of%20Haryana/2013-14.pdf> as accessed on May22, 2014



- Gangolli, Leena V., Duggal, Ravi and Shukla, Abhay (2005). Historical review of health care policy making: *Review of health care in India*, Section 2, 21-40.
- Haryana Government Portal <http://haryana.gov.in/haryana%20state/haryana%20state.asp> as accessed on May 22, 2014
- International institute for Population Sciences & Macro International (2007) *National Family Health Survey (NFHS-3)*, 2005-06: India: Volume I. Mumbai: IIPS
- International institute for Population Sciences & Macro International (2007) *National Family Health Survey (NFHS-3)*, 2005-06: India: Volume II. Mumbai: IIPS
- Jat, T.R., Ng, N., & San, Sebastian M. (2011). Factors affecting the use of maternal health services in Madhya Pradesh state of India: a multilevel analysis. *International Journal for Equity in Health*, 10, 59
- Misra, Rajiv, Chatterjee, Racheal and Rao, Sujatha (2003). *India Health Report*, Oxford University Press.
- National Health Policy (2002) [http://www.iapsmgc.org/userfiles/National\\_Health\\_policy\\_2002-2.pdf](http://www.iapsmgc.org/userfiles/National_Health_policy_2002-2.pdf) as accessed Feb 2014
- National Rural Health Mission (NRHM) (2005-10). [http://mohfw.nic.in/NRHM/Documents/5\\_Years\\_NRHM\\_Final.pdf](http://mohfw.nic.in/NRHM/Documents/5_Years_NRHM_Final.pdf) as accessed on March 10, 2014
- National Rural Health Mission Haryana [http://nrhm.gov.in/nrhm-in-state/state-wise-information/haryana.html#health\\_profileand](http://nrhm.gov.in/nrhm-in-state/state-wise-information/haryana.html#health_profileand) Potential: *Medical care*, 16(7), 533-546
- National Rural Health Mission Haryana <http://www.nrhmharyana.gov.in/programDetails.aspx?id=1> as accessed on 5<sup>th</sup> June, 2014
- Navaneetham, K., & Dharmalingam, A. (2002). Utilization of maternal health care services in southern India. *Social Science and Medicine*, 55(10), 1849-1869.
- Pallikadavath, S., Foss, M., & Stones, R.W. (2004). Antenatal care: provision and inequality in rural north India. *Social Science and Medicine*, 59(6), 1147-1158.

Pandey, A., Roy, N., Sahu, D. and Acharya, R. (2004). Maternal health care services: Observations from Chhattisgarh, Jharkhand and Uttaranchal. *Economic and Political Weekly*, Feb 14 ,713-720.

Saroha, E., Altarac, M., & Sibley, L.M. (2008). Caste and maternal health care service use among rural Hindu women in Maitha, Uttar Pradesh, India. *Journal of midwifery & Women's Health*, 53(5), e41-e47

STATA 10 Statistical package for Data Analysis- STATA corp

Sunil, T.S., Rajaram, S., Zottarelli, L.K. (2006). Do individual and program factors matter in the utilization of maternal care services in rural India? A theoretical approach. *Social Science and Medicine*, 62(8), 1943-1957.

World Health Organization. (2014).

<http://www.who.int/mediacentre/factsheets/fs290/en/> as accessed Jan 2015